arXiv:1212.6424v1 [nlin.SI] 27 Dec 2012

FACTORIZATION PROBLEM WITH INTERSECTION

R. A. ATNAGULOVA, O. V. SOKOLOVA

Abstract. We propose a generalization of the factorization method to the case when $\mathcal{G}$ is a finite dimensional Lie algebra such that $\mathcal{G} = \mathcal{G}_0 \oplus M \oplus N$ (direct sum of vector spaces), where $\mathcal{G}_0$ is a subalgebra in $\mathcal{G}$, M, N are $\mathcal{G}_0$-modules, and $\mathcal{G}_0 + M$, $\mathcal{G}_0 + N$ are subalgebras in $\mathcal{G}$. In particular, we consider the case when $\mathcal{G}$ is a $\mathbb{Z}$-graded Lie algebra. Using this generalization, we construct some top-like systems related to the algebra $so(3,1)$. According to the general scheme, these systems can be reduced to linear systems with variable coefficients. For the top-like systems first integrals and infinitesimal symmetries are found.

Key words: factorization method, Lie algebra, integrable dynamical systems.

1. Introduction

The classical factorization method [1, 2, 3, 4] (another name is Adler-Konstant-Symes scheme) allows to integrate the ODE system of the following special form:

$$U_t = [U_+, U], \quad U(0) = U_0. \tag{1}$$

Here U_t is a $\mathcal{G}$-valued function, where $\mathcal{G}$ is a Lie algebra, which is a direct sum of vector spaces $\mathcal{G}_+$ and $\mathcal{G}_-$ being subalgebras in $\mathcal{G}$. By U_+ we denote the projection of U onto $\mathcal{G}_+$. For simplicity we assume that $\mathcal{G}$ is embedded into the matrix algebra.

The solution of problem (1) is defined by the formula

$$U(t) = A(t)U_0A^{-1}(t). \tag{2}$$

The matrix $A(t)$ in (2) is defined as a solution of the factorization problem

$$A^{-1}B = \exp(-U_0t), \quad A \in G_+, \quad B \in G_-, \tag{3}$$

where G_+ and G_- are the Lie groups of the algebras $\mathcal{G}_+$ and $\mathcal{G}_-$, correspondingly. If $\mathcal{G}_-$ is an ideal, then the factorization problem can be solved explicitly: $A = \exp((U_0)_+t)$, $B = A\exp(-U_0t)$. In the case when G_+ and G_- are algebraic groups, the conditions $A \in G_+$ and $A\exp(-U_0t) \in G_-$ yield a system of algebraic equations. From this system the matrix A for small t is uniquely defined.

It was shown in [4] that the classical factorization problem can be reduced to a system of ordinary differential equations with variable coefficients.

Moreover, in [4] the factorization method was generalized to the case

$$\mathcal{G} = V_1 \oplus V_2, \tag{4}$$

where V_1, V_2 are vector subspaces in $\mathcal{G}_+$ and $\mathcal{G}_-$, correspondingly. It was shown that if

$$[\mathcal{G}_+ \cap \mathcal{G}_-,\, V_i] \subset V_i, \qquad i = 1, 2, \tag{5}$$

then the integration of equation (1), where «+» means the projection onto V_1 parallel to V_2, can be also reduced to solving of a system of ODEs with variable coefficients.

Recall the Golubchik-Sokolov construction. In [4] the authors have considered a factorization problem with *right* logariphmic derivatives of A and B:

$$A^{-1}B = Z(t), \quad A_tA^{-1} \in V_1, \quad B_tB^{-1} \in V_2, \quad Z(0) = A(0) = B(0) = E, \tag{6}$$

where V_i are vector spaces satifying (4). Let $U(t) = A(t)q(t)A^{-1}(t)$, where $q(t) = -Z_tZ^{-1}$. It is easy to see that this function satisfies the following equation:

$$U_t = [U_+, U] + Aq_tA^{-1}. \tag{7}$$

Notice that if $Z(t) = \exp(-U_0t)$ then $q(t) = U_0$, $q_t = 0$ and equation (7) coincides with (1). Thus, any solution of this factorization problem yields a solution of non-linear ODE (1).

Consider now the following factorization problem:

$$\alpha^{-1}\beta = Z(t), \quad \alpha^{-1}\alpha_t \in V_1, \quad \beta^{-1}\beta_t \in V_2, \quad Z(0) = \alpha(0) = \beta(0) = E. \tag{8}$$

Here the both logariphmic derivatives are *left*. In general, this problem does not related to equations of the (1) type. But it can be reduced to a linear equation with variable coefficients. Namely, in [4] have been introduced the linear mapping $L(t) : V_1 \to V_1$, defined by

$$L(t)(v) = (Z^{-1}(t)vZ(t))_+.$$

Since $L(0)$ is the identity mapping, $L(t)$ is invertible for small t. It can be proved that the solution α of the linear equation

$$\alpha_t = -\alpha L^{-1}(t)((Z^{-1}Z_t)_+), \qquad \alpha(0) = E \tag{9}$$

and the function

$$\beta = \alpha Z(t), \tag{10}$$

define the unique solution of the factorization problem (8).

The last step of the construction allows us to establish a link between the two factorization problems under additional condition (5). Namely, let $V_1 \subset \mathcal{G}_+$, $V_2 \subset \mathcal{G}_-$, where $\mathcal{G}_+$ and $\mathcal{G}_-$ are Lie subalgebras of the algebra $\mathcal{G}$ such that $\mathcal{G}_+ \cap \mathcal{G}_- = \mathcal{G}_0 \neq \{0\}$. Then the solutions of the factorization problems (6) and (8) satisfy the same factorization problem:

$$A^{-1}B = Z(t), \quad A \in G_+, \quad B \in G_-, \quad A(0) = B(0) = E, \tag{11}$$

where G_+ and G_- are the Lie groups of the algebras $\mathcal{G}_+$ and $\mathcal{G}_-$. Since $\mathcal{G}_0 \neq \{0\}$, the solution of problem (11) is not unique. Let α, β be the solution of the factorization problem (8). Because it also satisfies (11), all solutions of (11) are given by

$$A = H\alpha, \quad B = H\beta, \quad H(0) = E, \tag{12}$$

where H is an arbitrary element of the Lie group G_0 of the Lie algebra $\mathcal{G}_0$.

In [4] it was shown that A, B satisfy the factorization problem (6) if H is the solution of the following linear equation:

$$H_t = -H((\alpha_t\alpha^{-1})_- + (\beta_t\beta^{-1})_+), \quad H(0) = E. \tag{13}$$

Thus we solve equation (9) for α, find function β from (10) and solving equation (13), we determine the solution A, B of (6) from (12). Now we can find the solution of equation (7) by formula (2).

In this paper we generalize the factorization method to the case when $\mathcal{G}$ is a finite dimensional Lie algebra such that $\mathcal{G} = \mathcal{G}_0 \oplus M \oplus N$ (direct sum of vector spaces), where $\mathcal{G}_0$ is a subalgebra in $\mathcal{G}$, M, N are $\mathcal{G}_0$-modules, and $\mathcal{G}_0 + M$, $\mathcal{G}_0 + N$ are subalgebras in $\mathcal{G}$. The graded Lie algebras provide an important special class for which our generalisation is applicable. We construct some top-like systems related to the algebra $so(3,1)$. According to the general scheme, these systems can be reduced to linear systems with variable coefficients. For the top-like systems first integrals and infinitesimal symmetries are found.

2. General construction

Suppose we have a decomposition

$$\mathcal{G} = \mathcal{G}_0 \oplus M \oplus N$$

of a finite dimensional Lie algebra $\mathcal{G}$ over $\mathbb{R}$ into a vector space direct sum of a Lie subalgebra $\mathcal{G}_0$ and two vector spaces M, N such that

- M, N are $\mathcal{G}_0$-modules;
- $\mathcal{G}_0 + M$, $\mathcal{G}_0 + N$ are subalgebras in $\mathcal{G}$.

Theorem. *Let a linear operator R is defined by the formula*

$$R(q)=\alpha_{-1}q^{-1}+\alpha_0 q^0+\alpha_1 q^1, \tag{14}$$

where $q=q^{-1}+q^0+q^1$, $q^{-1}\in N$, $q^0\in\mathcal{G}_0$, $q^1\in M$, $\alpha_{-1},\alpha_0,\alpha_1\in\mathbb{R}$. Then the equation

$$q_t=[R(q),q], \quad q|_{t=0}=q_0, \tag{15}$$

can be reduced to a system of linear ODEs with variable coefficients by the Golubchik-Sokolov construction (see Introduction).

Proof. To apply the construction from [4], we take for $\bar{\mathcal{G}}$ the direct sum of three copies of the algebra $\mathcal{G}$, i.e.

$$\bar{\mathcal{G}}=\mathcal{G}\oplus\mathcal{G}\oplus\mathcal{G}.$$

For $\bar{\mathcal{G}}_+$ we take the diagonal subalgebra in $\bar{\mathcal{G}}$:

$$\bar{\mathcal{G}}_+=\{(a,a,a)|\, a\in\mathcal{G}\},$$

and for $\bar{\mathcal{G}}_-$ the following subalgebra:

$$\bar{\mathcal{G}}_-=\{(a,b,c)|\, a\in\mathcal{G}_0+N, b\in\mathcal{G}_0+M, c\in\mathcal{G}\}.$$

Consider the vector subspace

$$\bar{M}=\{(a,b,c)|\, a\in\mathcal{G}_0+N, b\in\mathcal{G}_0+M, c\in N+M\}.$$

Then condition (4) holds for $\bar{\mathcal{G}}$ and $\bar{\mathcal{G}}_+,\bar{M}$, i.e. $\bar{\mathcal{G}}=\bar{\mathcal{G}}_+\oplus\bar{M}$. Obviously, for the subalgebra $\bar{\mathcal{G}}_+$ condition (5) is fulfilled. It is easy to verify that for $\bar{M}$ we have

$$[(\bar{\mathcal{G}}_+\cap\bar{\mathcal{G}}_-),\bar{M}]\subset\bar{M}.$$

Thus it follows from [4] that the equation

$$\bar{q}_t=[\bar{q}_+,\bar{q}], \quad \bar{q}|_{t=0}=\bar{q}_0, \tag{16}$$

where $\bar{q}\in\bar{\mathcal{G}}$, $\bar{q}_+$ is the projection $\bar{q}$ onto $\bar{\mathcal{G}}_+$ parallel to $\bar{M}$, can be reduced to a system of linear ODEs with variable coefficients.

To complete the proof it is sufficient to choose $\bar{q}=(\alpha_1 q,\alpha_{-1}q,\alpha_0 q)$ where $q\in\mathcal{G}$. Notice, that in this case $\bar{q}_+=(R(q),R(q),R(q))$ where the operator R is defined by (14). Now one can see that for each component, equation (16) has the same form $q_t=[R(q),q]$. This coincides with (15). □

Corollary 1. *Let $\mathcal{G}=\bigoplus_{i=-k}^{k}\mathcal{G}_i$ be a $\mathbb{Z}$-graded Lie algebra and $N=\bigoplus_{i=-k}^{-1}\mathcal{G}_i$, $M=\bigoplus_{i=1}^{k}\mathcal{G}_i$. Then all conditions of Theorem are fulfilled and the equation*

$$q_t=\left[\alpha_{-1}\sum_{i=-k}^{-1}q_i+\alpha_0 q_0+\alpha_1\sum_{i=1}^{k}q_i,\ q\right] \tag{17}$$

can be reduced to a system of linear equations with variable coefficients.

3. Examples

Corollary gives rise to the following two interesting examples.

Example 1. Consider the following graduation on $\mathcal{G}=\mathfrak{sl}_2$:

$$\mathcal{G}_{-1}=\left\{\begin{pmatrix}0&0\\v&0\end{pmatrix}\right\}, \quad \mathcal{G}_0=\left\{\begin{pmatrix}w&0\\0&-w\end{pmatrix}\right\}, \quad \mathcal{G}_1=\left\{\begin{pmatrix}0&u\\0&0\end{pmatrix}\right\}.$$

The equation (17) can be rewritten as the system

$$\begin{cases} w_t &= -(\alpha_1-\alpha_{-1})\,uv,\\ u_t &= 2\alpha_0\,uw,\\ v_t &= 2\alpha_0\,vw. \end{cases}$$

The system possesses expected first integral

$$H_1 = \frac{1}{2} Tr(q^2) = uv + w^2$$

and, in addition, it has the following transcendental first integral

$$H_2 = u^{\alpha_0-\alpha_{-1}} v^{\alpha_0-\alpha_1}.$$

Integrable dynamical systems with quadratic right hand sides associated with semi-simple Lie algebras have been considered in [5, 6, 7]. The complex isomorphic algebras $so(4)$ and $so(3,1)$ are interesting from a viewpoint of applications. Integrable Hamiltonian systems associated with these algebras were studied in [8, 9, 10]. We present two new examples related to the algebra $so(3,1)$. Systems from these examples apparently are non-Hamiltonian. Furthermore, their first integrals just as in Example 1 are not single valued in the complex plane. However, since the systems are linearizable, they should be regarded as integrable in some sense. We produce sufficiently many first integrals and infinitesimal symmetries to claim that these systems are locally integrable by the theorem of S. Lie.

Example 2. Let $\mathcal{G} = \{A \in \mathbb{R}_{4\times 4} |\, A^* = -A\}$ where the involution is defined by the formula $A^* = TA^tT^{-1}$, $T = e_{11} + e_{22} + e_{34} + e_{43}$, the superscript t means the matrix transposition, and e_{ij} denote the matrix unities. If we take $T = e_{11} + e_{22} + e_{33} - e_{44}$ then the condition of skew-symmetricity with respect to the involution defines the algebra $so(3,1)$. The isomorphism between $\mathcal{G}$ and $so(3,1)$ is given by $\bar{q} = B^{-1}\, q\, B$, where $q \in \mathcal{G}$, $\bar{q} \in so(3,1)$, and the matrix B equals

$$e_{11} + e_{22} + \frac{\sqrt{2}}{2}(e_{33} + e_{34} + e_{43} - e_{44}).$$

Define a 3-graduation on the algebra $\mathcal{G}$. Generic element of the algebra has the form

$$q = \begin{pmatrix} 0 & w_1 & u_1 & v_1 \\ -w_1 & 0 & u_2 & v_2 \\ -v_1 & -v_2 & w_2 & 0 \\ -u_1 & -u_2 & 0 & -w_2 \end{pmatrix}.$$

It is easily verified that the decomposition of $\mathcal{G}$ into the sum of the following components

$$\begin{aligned} \mathcal{G}_0 &= \{w_1(e_{12} - e_{21}) + w_2(e_{33} - e_{44})\}, \\ \mathcal{G}_1 &= \{u_1(e_{13} - e_{41}) + u_2(e_{23} - e_{42})\}, \\ \mathcal{G}_{-1} &= \{v_1(e_{14} - e_{31}) + v_2(e_{24} - e_{32})\}. \end{aligned}$$

equips $\mathcal{G}$ with a structure of a 3-graded Lie algebra.

In this case equation (17) can be written as the system

$$\begin{cases} w_{1t} &= (\alpha_1 - \alpha_{-1})\,(u_2v_1 - u_1v_2), \\ u_{1t} &= (\alpha_0 - \alpha_1)\,(u_2w_1 - u_1w_2), \\ v_{1t} &= (\alpha_0 - \alpha_{-1})\,(v_2w_1 + v_1w_2), \\ w_{2t} &= (\alpha_1 - \alpha_{-1})\,(u_1v_1 + u_2v_2), \\ u_{2t} &= (\alpha_1 - \alpha_0)\,(u_1w_1 + u_2w_2), \\ v_{2t} &= (\alpha_{-1} - \alpha_0)\,(v_1w_1 - v_2w_2). \end{cases}$$

The system possesses the following quadratic infinitesimal symmetry:

$$\begin{cases} w_{1\tau} &= (-\alpha_1 + \alpha_{-1})\,(u_1v_1 + u_2v_2), \\ u_{1\tau} &= (-\alpha_0 + \alpha_1)\,(u_1w_1 + u_2w_2), \\ v_{1\tau} &= (\alpha_0 - \alpha_{-1})\,(v_1w_1 - v_2w_2), \\ w_{2\tau} &= (-\alpha_1 + \alpha_{-1})\,(u_1v_2 - u_2v_1), \\ u_{2\tau} &= (\alpha_1 - \alpha_0)\,(u_2w_1 - u_1w_2), \\ v_{2\tau} &= (\alpha_0 - \alpha_{-1})\,(v_2w_1 + v_1w_2). \end{cases}$$

This system has two polynomial first integrals of degree two:

$$I_1 = w_1w_2 + u_1v_2 - v_1u_2$$

and

$$I_2 = w_1^2 + 2u_1v_1 - w_2^2 + 2u_2v_2.$$

The traces of powers of q are polynomials in these two integrals.

Moreover, the system possesses first integrals of the form $I = f(u_1, u_2, v_1, v_2)$. It is easy to check that such integrals should satisfy two equations of the form $X(f) = 0$ and $Y(f) = 0$, where X, Y are given by:

$$X = -(\alpha_0 - \alpha_1)u_1\frac{\partial}{\partial u_2} - (\alpha_0 - \alpha_{-1})v_1\frac{\partial}{\partial v_2} + (\alpha_0 - \alpha_1)u_2\frac{\partial}{\partial u_1} + (\alpha_0 - \alpha_{-1})v_2\frac{\partial}{\partial v_1},$$

$$Y = (\alpha_0 - \alpha_{-1})v_2\frac{\partial}{\partial v_2} - (\alpha_0 - \alpha_1)u_2\frac{\partial}{\partial u_2} + (\alpha_0 - \alpha_{-1})v_1\frac{\partial}{\partial v_1} - (\alpha_0 - \alpha_1)u_1\frac{\partial}{\partial u_1}.$$

Solving this system of two PDEs, we find it's simplest solutions

$$I_3 = \operatorname{tg}\left(\frac{\alpha_{-1} - \alpha_0}{\alpha_1 - \alpha_0}\operatorname{arctg}\frac{u_1}{u_2} - \operatorname{arctg}\frac{v_1}{v_2}\right),$$

$$I_4 = (u_1^2 + u_2^2)^{\frac{\alpha_{-1}-\alpha_0}{\alpha_1-\alpha_0}}(v_1^2 + v_2^2)\sin^2\left(\frac{\alpha_{-1} - \alpha_0}{\alpha_1 - \alpha_0}\operatorname{arctg}\frac{u_2}{u_1} - \operatorname{arctg}\frac{v_2}{v_1}\right).$$

In the next example the algebra $\mathcal{G}$ is not $\mathbb{Z}$-graded.

Example 3. Let $\mathcal{G}$ be the same algebra as in Example 2. Now we consider

$$\mathcal{G}_0 = \left\{\begin{pmatrix} 0 & w_1 & 0 & 0 \\ -w_1 & 0 & 0 & 0 \\ 0 & 0 & kw_1 & 0 \\ 0 & 0 & 0 & -kw_1 \end{pmatrix}\right\},$$

$$M = \left\{\begin{pmatrix} 0 & lw_2 & 0 & v_1 \\ -lw_2 & 0 & 0 & v_2 \\ -v_1 & -v_2 & w_2 & 0 \\ 0 & 0 & 0 & -w_2 \end{pmatrix}\right\},$$

$$N = \left\{\begin{pmatrix} 0 & 0 & u_1 & 0 \\ 0 & 0 & u_2 & 0 \\ 0 & 0 & 0 & 0 \\ -u_1 & -u_2 & 0 & 0 \end{pmatrix}\right\},$$

where k and l are parameters such that $kl \neq 1$. It is easy to check that all conditions of the Theorem hold. Equation (15) can be rewritten in the form:

$$\begin{cases} w_{1t} &= \dfrac{\alpha_1 - \alpha_{-1}}{1 - kl}(-lu_1v_1 + u_2v_1 - u_1v_2 - lu_2v_2), \\ u_{1t} &= (\alpha_1 - \alpha_0)(ku_1 - u_2)w_1 + (\alpha_1 - \alpha_{-1})(u_1 - lu_2)w_2, \\ v_{1t} &= (\alpha_0 - \alpha_{-1})(kv_1 + v_2)w_1, \\ w_{2t} &= \dfrac{\alpha_1 - \alpha_{-1}}{1 - kl}(u_1v_1 - ku_2v_1 + ku_1v_2 + u_2v_2), \\ u_{2t} &= (\alpha_1 - \alpha_0)(u_1 + ku_2)w_1 + (\alpha_1 - \alpha_{-1})(lu_1 + u_2)w_2, \\ v_{2t} &= (\alpha_{-1} - \alpha_0)(v_1 - kv_2)w_1. \end{cases}$$

The trace of q^2 provides the following first integral:

$$I_1 = -2u_1v_1 - 2u_2v_2 + (kw_1 + w_2)^2 - (w_1 + lw_2)^2.$$

Acknowledgements.

The authors are deeply grateful to I. Z. Golubchik and V. V. Sokolov for the useful discussions and the attention to the work. The research of O. S. was partially supported by the RFBR grant 11-01-00341-a.

References

[1] B. Konstant. *Quantization and representation theory.* Lect. Notes Ser., 1979, 34, p. 287–316.
[2] M. A. Semenov-Tyan-Shansky. *What is a classical r-matrix?* (in Russian) // Func. Anal. and Appl., 1983, **7**, №4, p. 17—33.
[3] P. I. Golod. *Hamiltonian systems on orbits of affine Lie groups and nonlinear integrable systems* (in Russian) // In Fizika mnogochastichnyh sistem, Kiev, 1985, **7**, p. 30–39.
[4] I. Z. Golubchik, V. V. Sokolov. *On some generalizations of the factorization method.* Teoret. and Mat. Fiz, 1997, **110**, №3, p. 339-350
[5] A. S. Mitschenko, A. T. Fomenko. *Euler equations on finite dimensional Lie groups* (in Russian) // Isvestia AN SSSR, math., 1978, **42**, №2, p. 396 – 415.
[6] A. G. Reyman, M. A. Semenov-Tyan-Shansky. *Integrable systems. Group theory approach* (in Russian) // Izhevsk, 2003, 351 p.
[7] A. V. Borisov, I. S. Mamaev. *Solid body dynamics* (in Russian) // Izhevsk, 2001, 384 p.
[8] A. P. Veselov. *On integrability conditions of Euler equation on* $so(4)$ (in Russian) // DAN SSSR, 1983, 270, №6, p. 1298–1300.
[9] V. V. Sokolov. *One class of quadratic* $so(4)$*-Hamiltonians* (in Russian) // Doklady RAN, 2004, 394(5), p. 602–605. Translation in *Doklady Mathematics*, 2004, 69, №1, p. 108–111.
[10] V. V. Sokolov, T. Wolf. *New integrable quadratic Hamiltonians on* $so(4)$ *and* $so(3,1)$, Journal Phys. A: Math. Gen., 2006, 39, p. 1915-1936.